



Purpose

Asthma affects up to one in four primary aged children, one in seven teenagers and one in ten adults. It is important therefore for all staff members to be aware of asthma, its symptoms and triggers, and the management of asthma in a school environment.

Scope

To manage asthma and asthma sufferers as effectively and efficiently as possible at school.

Policy

Asthma attacks involve the narrowing of airways making it difficult to breathe. Symptoms commonly include difficulty breathing, wheezy breathing, dry and irritating cough, tightness in the chest and difficulty speaking.

- Children and adults with mild asthma rarely require medication, however severe asthma sufferers may require daily or additional medication (particularly after exercise).
- Professional development will be provided annually for all staff on the nature, prevention and treatment of asthma attacks. Such information will also be displayed in the first aid room.
- All students with asthma must have an up to date (annual) written asthma management plan consistent with Asthma Victoria's requirements completed by their doctor or paediatrician. Appropriate asthma plan proformas are available at www.asthma.org.au
- Asthma plans will be attached to the student's records for reference.
- Parents/guardians are responsible for ensuring their children have an adequate supply of appropriate asthma medication (including a spacer) with them at school at all times.
- The school will provide, and have staff trained in the administering of, reliever puffers (blue canister) such as Ventolin, Airomir, Asmol or Bricanyl and spacer devices in all first-aid kits, including kits on excursions and camps.
- The first aid staff member will be responsible for checking reliever puffer expiry dates.
- Care must be provided immediately for any student who develops signs of an asthma attack.
- Children suffering asthma attacks should be treated in accordance with their asthma plan.
- If no plan is available children are to be sat down, reassured, administered 4 puffs of a shaken reliever puffer (blue canister) delivered via a spacer – inhaling 4 deep breaths per puff, wait 4 minutes, if necessary administer 4 more puffs and repeat the cycle. An ambulance must be called if there is no improvement after the second 4-minute wait period, or if it is the child's first known attack. Parents must be contacted whenever their child suffers an asthma attack.
- The school will purchase Ventolin puffers for general use. These will be stored in the first aid room and in backpacks which will be taken on sports days, excursions, camps and special events.
- All yard duty bags will contain Asthma Action Kits.

Policy review

Policy written updated	School Council approved	Review date	Responsibility
February 2021	N/A	February 2022	First Aid Coordinator